

## A New Species of *Alpinia* (Zingiberaceae) from Vietnam

NGUYEN QUOC BINH

Institute of Ecology and Biological Resources NCST,  
Nghia Do, Cau Giay, Hanoi, Vietnam

AND

MARK NEWMAN

Royal Botanic Garden, Edinburgh EH3 5LR, U.K.

### Abstract

*Alpinia calcicola* Q.B. Nguyen & M.F. Newman, a new species of subsect. *Alpinia* from Ha Long Bay, Vietnam, is described.

### Introduction

While working on a guide to plants of Ha Long Bay (N.T. Nguyen & Kiew, 2000), a species of *Alpinia* was collected that does not match any of those cited in Q.B. Nguyen (1994). Further study indicates that it is a new species, which is described here:

***Alpinia calcicola* Q.B. Nguyen & M.F. Newman sp. nov.** *Alpiniam maclurei* Merr. tangit, ob laminam minorem, (22—31 x 3—7 cm), glabram, bracteolam 2 vel 3 flores gerentem, tubum calycis tubum corollae aequantem, antheram ecristatam inter species subsectionis *Alpiniae* ponenda.

**Typus:** Vietnam, Quang Ninh Province, Ha Long Bay, *Nguyen Tien Hiep & Ruth Kiew, NTH 4220* 12 April 2000, flowering (holo HN; iso SING).

Herb 1.5—2 m. *Leaf sheath* glabrous, striate; ligule 8—10 mm, emarginate, shortly hairy, ciliate at margin; *petiole* 2—7 mm, reddish brown, glabrous; *lamina* lanceolate to linear-lanceolate, 22—31 x 3—7 cm, light green above, yellowish green beneath with reddish brown midrib, glabrous on both surfaces; margin with sparse, tooth-like hairs; base acute; apex acuminate. *Inflorescence* terminal, yellowish brown, 26—32 cm long; rachis slender, 2—3 mm diameter, sparsely hairy; *cincinni* diffuse, becoming denser towards apex, axes hairy; bract subtending inflorescence pale pinkish brown, to 18 x 2 cm, caducous; *cincinni* without bracts, usually with a very short flap of free tissue at the join with the rachis; *bracteoles* broadly ovate, split to the

base, pinkish white to pink, hairy at tip and margin, usually subtending 2—3 flowers, first bracteole 12—14 x 8—10 mm, subsequent ones smaller; *ovary* globose, hirsute; epigynous glands *c.* 3 mm long, fused into a truncated cone; *calyx* tubular, 12—14 mm, adpressed silky hairy outside, split into three short ciliate teeth above; *corolla* tube as long as calyx, very sparsely hairy outside, dorsal corolla lobe 14—19 x 4—6 mm, oblong, slightly hooded, lateral corolla lobes smaller, 14—16 x 5—7 mm; labellum *c.* 14 x 18 mm, broader than long, shortly bifid at apex, margins frilled, lateral lobes pure crimson with white margin, centre with longitudinal crimson lines; *lateral staminodes* 4—5 mm, fleshy; filament 13—14 mm, anther 4—5 mm long, sparsely short hairy on back, thecae parallel, connective sometimes produced as a small flap of tissue; *style* filiform, *stigma* funnel-form, ciliate; *fruit* globose, 6—7 mm.

*Other specimens seen:* Vietnam, Quang Ninh Province, Ha Long Bay: Nguyen Tien Hiep & Leonid Averyanov NTH2600 12 May 1999 (HN, LE, SING); Nguyen Tien Hiep & Ruth Kiew NTH3931 15 July 1999 (HN, SING).

*Habitat:* In rock cracks and crevices in open places from sea level almost to the summits of limestone islands.

*Notes:* This species is illustrated in colour by Wendy Gibbs in Plate 17 of *Wild Plants of Ha Long Bay* (T.N. Nguyen and Kiew, 2000).

### Acknowledgements

We should like to thank Nguyen Tien Hiep (HN), Ruth Kiew (SING) and Leonid Averyanov (LE) for bringing this new species to our attention and providing us with material. This species was discovered during the fieldwork for the guide to Wild Plants of Ha Long Bay, a project initiated and coordinated by IUCN Vietnam.

### References

- Nguyen, Q.B. 1994. The genera *Alpinia* Roxb. and *Kaempferia* L. (Zingiberaceae) in the flora of Vietnam. *Journal of Biology (Viet Nam)* 16(4): 143–145.
- Nguyen, T. H. & R. Kiew. 2000. *Wild Plants of Ha Long Bay*. IUCN Vietnam, Hanoi, Vietnam. Illustrated by W. Gibbs.